

# EXPERIMENTAL ASTROPARTICLES 1

24 July 2017

**Vincent Poireau, LAPP Annecy**

# PLAN


- Experimental astroparticles 1
  - Cosmic rays
  - Indirect search for dark matter
  - Some experiments
  - AMS-02: detailing a modern experiment
  - Recent results on cosmic rays and their implications
- Experimental astroparticles 2
  - This afternoon, presented by **David Maurin**
  - The gamma-ray sky
  - Interactions in a detector/atmosphere
  - Ground vs space detectors
  - Fermi-LAT and H.E.S.S. (and Auger)
  - Recent results and constraints on dark matter

# COSMIC RAYS



# HISTORIC

- 1736 – 1806 : **Charles Augustin de Coulomb** observed that a sphere initially charged and isolated loses its electrical charge




# HISTORIC

- 1736 – 1806 : **Charles Augustin de Coulomb** observed that a sphere initially charged and isolated loses its electrical charge



Electroscopes  
are  
spontaneously  
discharging  
????




# HISTORIC

???

Space



Beginning of 20<sup>th</sup> century



Henri Becquerel  
(1852-1908)

Marie Curie  
(1867-1934)

*The radioactivity  
could explain  
the spontaneous discharge*






C.T.R. Wilson  
(1869-1959)

Ground



# HISTORIC



- 1912: **Victor Hess** measures the atmospheric ionization with electroscopes during balloon flights at various altitudes: the **ionization increases**



- This ionization comes from **space!**

# HISTORIC

- From what are they **composed**? The debate is **passionate** in the 1920's



1925: very high energy  
gammas → « cosmic rays »



- Their intensity **varies** depending on where we are on Earth...
- Cosmic rays are **charged particles**!
  - More particle from the western direction: **positively charged**

# HISTORIC

- 1937: **Pierre Auger** positions three Geiger counters separated of 70 m at le pic du midi
- Cosmic rays arrive in group: **atmospheric shower**





Pierre Victor Auger  
(1899-1993)





# HISTORIC

- Many new particles discovered in the cosmic rays
  - 1932: positron  $e^+$  (first observation of antimatter)
  - 1936: muon  $\mu$
  - 1949: pion  $\pi$
  - 1949: kaon  $K$
  - 1949: lambda  $\Lambda$
  - 1952: xi  $\Xi$
  - 1953: sigma  $\Sigma$
- Birth of a new science: particle physics!
- Cosmic rays are replaced by accelerators where particles are artificially produced





# COSMIC RAYS

- **Cosmic rays**
  - 12 orders in **energy**
    - 100 MeV to  $10^{20}$  eV
  - 30 orders in **flux**
  - Isotropic flux
- **Abundance** of nuclei in the cosmic rays similar to the one from the **solar system**



# COSMIC RAYS


- **Composition**
  - **Charged** : electrons, protons, nuclei
  - **Neutral** : photons, neutrinos
- **Charged** cosmic rays




- **Power law spectrum**  $1/E^\gamma$ ,  $\gamma = 2.7-3.5$ 
  - The measured spectrum results
    - from the **production** and **acceleration** mechanisms ( $1/E^\alpha$ ,  $\alpha = 2.0-2.4$ )
    - from the **diffusion** ( $1/E^\delta$ ,  $\delta = 0.3-0.7$ )
  - $\gamma = \alpha + \delta$

# COSMIC RAYS

- Where are they coming from?



From the sun




From outside our Galaxy

From our Galaxy



# COSMIC RAYS

- Low energy cosmic rays are accelerated by **the sun**



Aurora borealis




# COSMIC RAYS

- At intermediate energies, **supernovae remnants** produce cosmic rays




# COSMIC RAYS

- At extreme energies, active galaxy nuclei, quasars, or gamma ray bursts are potential candidates



Centaurus A


# COSMIC RAYS



- **Primary cosmic rays**
  - Produced directly **in the source**
  - Sources: **supernova remnants, pulsars, active galactic nuclei, quasars**
  - Primaries include
    - Electrons, protons, helium, carbon, ...
- **Secondary cosmic rays**
  - Originate from the **interaction** of primaries on **interstellar medium**
  - Secondaries include
    - Positrons, antiprotons, boron, ...
- **Additional sources of electrons and positrons?**



# ACCELERATION

- In our Galaxy, **main source** of primary cosmic rays: **supernova remnants**
  - Very strong magnetic field in the **shell** of supernovas
- Acceleration
  - Due to the **shock wave**
  - First order **Fermi mechanism**
  - Naturally produce a **power law** spectrum
- This process explains why the cosmic ray **composition is similar** to the one from the solar system



# PROPAGATION



- Charged cosmic rays: propagation equivalent to a **diffusion** in the Galactic medium
  - **Irregular magnetic field** of the diffusive halo = random walk
  - Possible to write a **cosmic ray transport equation**
    - Difficult to solve
    - **5 free parameters**, with large **uncertainties** on these parameters




$h_z=200$  pc,  $L=1-15$  kpc,  $R=25$  kpc

# SOLAR MODULATION

- **Heliosphere:** a region of space influenced by the sun (solar wind)
  - **Size:** 150 AU
- **Solar wind:** a continuous flow of charged particles from sun
  - $e^-$  and  $p$
  - Carries the **sun magnetic field** to the interplanetary space
- **Solar cycles**
  - **Reversal** of the sun magnetic field polarity
  - Every **11 years**
  - Solar activity going from a minimum to a maximal intensity
- **Solar modulation** affects cosmic rays below 20 GeV
  - **Deviation** from the power law




# INDIRECT SEARCH FOR DARK MATTER



# DARK MATTER


## Galaxy rotation curve



A

A : prediction


B : observation



To explain the observation, we have to suppose the presence of a halo containing an invisible matter around the galaxies: **the dark matter!**

# DARK MATTER


- A very large fraction of the Universe content remains mysterious



- Dark matter: 27% of our Universe is made of **unknown matter** (other than electrons, quarks, ...)
- « **Observation** »: galaxy rotation curves, X-ray emission, gravitational lensing, cosmic microwave background

# DARK MATTER


- **Best candidate:** weakly interacting massive particle  $\Rightarrow$  WIMP
  - **Massive particles:** 100 GeV – several TeV
  - **Weakly** interacting with the ordinary matter
- Several ways to see its effect



# DARK MATTER

- **Annihilation** of the WIMPs

- Natural cross-section from relic density:  $\langle\sigma v\rangle \approx 3 \cdot 10^{-26} \text{ cm}^3\text{s}^{-1}$



# COSMIC RAY EXPERIMENTS



# EXPERIMENTS


1959-1974: Vulcano Ranch (USA)  
8km<sup>2</sup>




1998-2004: AGASA (Japan)  
100 km<sup>2</sup>



1980-1993: Fly eye (Utah)



2004- : Pierre-Auger observatory



# EXPERIMENTS

1947: inside a B 29  
Altitude : 10 km



1979 à 1995: ECHO  
Altitude: 17 km




1965: proton satellite



Orbit: 183-589 km

1993-1996:  
JACEE



Alt: 30 km

1998: Discovery  
Altitude: 400 km




AMS-01

# EXPERIMENTS



2011-: AMS  
Altitude : 400 km  
Let's detail this experiment!



# THE AMS-02 EXPERIMENT



# AMS-02

- A **particle detector** in space
  - Detect **charged** particles and **gamma** rays
  - From **100 MeV** to a **few TeV**



5m x 4m x 3m  
7.5 tons

# AMS-02

- **Launched** from Cap Canaveral on the 16<sup>th</sup> of May 2011
  - **Penultimate** American shuttle!





Spacevidcast Live  
STS-134 Mission Coverage

T

-00:00:12

Official Clock


SSME Thrust: 0.0%  
Altitude (ft): -23

Speed (mph): 0  
Range (nm): 0.0

Mach: 0.00  
G: 0.0

# AMS-02


- Installation on the **ISS** on the 19<sup>th</sup> of May 2011
  - Orbit at **400 km** altitude
  - One orbit every **90 minutes**



- Detect the cosmic rays **before they interact** in the atmosphere

# FLIGHT OPERATION


- Acquisition rate from 200 to 2000 Hz
- Continuous operation 7d/7 24h/24
- Acquisition
  - ~40 millions events a day
  - ~100 GB transferred every day
  - 35 TB of data every year
  - 200 TB of reconstructed data every year
- More than 100 billions of events recorded since May 2011
  - Much more than all the cosmic rays collected in the last 100 years
- Will operate at least until 2024



Transition radiation  
detector  
Identifies  $e^+$ ,  $e^-$

# DETECTOR


Time of flight  
 $Z, E$





Silicium tracker  
 $Z, P$





Magnet 0,14 T  
 $\pm Z$



Electromagnetic calorimeter  
 $E$  of  $e^+$ ,  $e^-$ ,  $\gamma$




Cherenkov detector  
 $Z, E$



# DETECTOR

- **Rigidity**

- $R = p/Z$
- Expressed in GV




A 369 GeV positron event

# COLLABORATION



## AMS: a U.S. DOE sponsored international collaboration

16 Countries, 60 Institutes and 600 Physicists



# AMS TOPICS

- Measurement of **cosmic ray fluxes**
  - Understand the cosmic ray **propagation** in our Galaxy
- Indirect search of **dark matter**
  - **Positrons** and **antiprotons** produced during its annihilation
- Search for primordial **antimatter**
  - **Anti-helium** relic of the Big-Bang or **anti-carbon** from anti-stars
- Surprises? **Strangelets?**



# ELECTRONS AND POSITRONS IN COSMIC RAYS

# POSITRON FRACTION


- **Positrons** : expected only as secondary
- Positron excess with respect to the secondary prediction = **source of primary positrons**

- **Positron fraction** 
$$F = \frac{\Phi_{e^+}}{\Phi_{e^+} + \Phi_{e^-}} = \frac{N_{e^+}}{N_{e^+} + N_{e^-}}$$
  - Allows to factorize the **acceptance** and efficiencies
  - **Simplify** the computation of systematic uncertainties

- **Challenges**
  - **100 times** more protons than electrons
  - **2000 times** more protons than positrons
  - ⇒ Need to divide number of protons by **10<sup>6</sup>**


# POSITRON FRACTION

- Result for the positron fraction **below 35 GeV**
  - Fraction begins to increase **above 10 GeV**
  - **Incompatible** with secondary positrons only
  - A **source of primary positrons** is needed!
    - **Nearby source** since positrons do not propagate more than a **few kpc**



# POSITRON FRACTION

- Fraction at high energy



- AMS: precision and energy never reached before

# FLUX MEASUREMENT

- Fluxes bring more information for the models than the fraction


- Obtaining the flux via

$$\frac{N}{A \times \epsilon_{Trig.} \times \epsilon_{sel.} \times T \times dE}$$

- $N$  **number** of positrons or electrons
- $A$  **acceptance**
- $\epsilon_{Trig}$  and  $\epsilon_{sel}$  **trigger and selection efficiencies**
- $T$  **exposure time**
- $dE$  **energy bin size**


# FLUX MEASUREMENT

- The electron and positron fluxes are **different** in their **magnitude** and **energy dependence**



# INTERPRETATION: DARK MATTER

- Fitting the positron fraction using the **best combination** of annihilation channels





A&A 575, A67 (2015)

- Dark matter **may explain** the fraction, but **unnatural** annihilation cross-section
  - $\times 1000$  compared to the one expected from the relic density
- Not likely** that we have observed an indirect observation of dark matter
  - In tension** with other observables (antiprotons, gamma rays, ...)

# INTERPRETATION: PULSARS

- Neutron stars spinning at high rate with a strong magnetic field
- 200 pulsars at less than 2 kpc from Earth
  - Only a **small fraction** able to emit positrons
- Mechanism
  - **Electrons extracted** from the surface by the high fields
    - ⇒ electrons produce **synchrotron photons**
    - ⇒ photons produce  **$e^+e^-$  pairs**
    - ⇒ Some **escape** from the pulsar
- Precise prediction **very difficult**
- Five closeby pulsars able to explain the fraction




A&A 575, A67 (2015)

# ANTIPROTONS IN COSMIC RAYS


# ANTIPROTONS

- Dark matter could create **an excess of antiprotons** with respect to the expectations
  - Pulsars **do NOT** produce antiprotons
- AMS measured the ratio  $\bar{p}/p$ 
  - 290 000 antiprotons
- Is dark matter **necessary** to explain this measurement?
  - Controversial topic!
  - Need to compute what is expected from **secondary antiprotons**



# ANTIPROTONS





- Adding the contribution of the **secondary antiprotons** with its uncertainty
  - **Comparison** of data and expectations for  $\bar{p}/p$



- The ratio  $\bar{p}/p$  is **not in discrepancy** with the expectations
  - **No dark matter** needed here (yet)


# OTHER RESULTS ON COSMIC RAYS

# OTHER RESULTS



# NUCLEUS ANOMALY

- Recent and **unexpected** observation: the power law is **broken** at high energy




- What can cause these **anomalies**?
  - Sources?
  - Acceleration?
  - Propagation?
- Still an **open question**!

# BORON/CARBON RATIO

- Ratio of secondary over primary allows to understand the **propagation** of cosmic rays
  - **Carbon**: primary, produced and accelerated in sources
  - **Boron**: secondary, produced by the collision of heavier nuclei on the interstellar matter
  - **Ratio** directly measures the average amount of interstellar material traversed by the cosmic rays

**Bonus:** a **break** in the B/C ratio would explain the nucleus anomaly by propagation phenomena



# IN SUMMARY

- **Cosmic rays** are charged and neutral particles coming from space
  - From a few MeV to  $10^{20}$  eV
  - Mainly **protons, helium, electrons, ...**
- **Sources**
  - At intermediate energies, they come from **supernova remnant** in our Galaxy
  - Protons, electrons, ... come directly **from the source**
  - Positrons, antiprotons, ... are created by **collision** with the interstellar medium, with a rate that **can be predicted**
- **Propagation**
  - Charged cosmic ray propagation is equivalent to a **diffusion**
- **Positrons in cosmic rays**
  - There is **more positrons** at high energy compared to the expectations
  - **New source**: dark matter? pulsars?
  - AMS will extend its **energy range**, and should be able to **discriminate** between the dark matter and pulsar hypotheses
- **Antiprotons in cosmic rays**
  - Antiprotons could be produced by dark matter
  - After the recent AMS measurement, **no need for dark matter**
- **Other measurements**
  - Many other measurement are **yet to come**, with on-going experiments or **promising future experiments** (CALET, DAMPE, ISS-CREAM)

TO BE CONTINUED...  
(David Maurin)

# ADDITIONAL SLIDES

# WIMP “miracle”

- Start with **heavy, stable dark matter** (DM) particle  $X$  in thermal equilibrium.
- **Early universe**  $T > M_X$ :  $X\bar{X} \leftrightarrow f\bar{f}$
- **Universe cools**  $T < M_X$ :  $X\bar{X} \rightarrow f\bar{f}$
- **Freeze out**: Hubble expansion eventually prevents  $X\bar{X} \rightarrow f\bar{f}$
- Solving Boltzmann equation


$$\frac{dn}{dt} = -\frac{3\dot{R}}{R} n - \langle\sigma v\rangle n^2 + \langle\sigma v\rangle n_0^2$$

assuming measured DM density results in:

$$\frac{\Omega_{\text{DM}} h^2}{0.1} \approx \left( \frac{\langle\sigma v\rangle}{3 \text{ pb} \cdot \text{cm/s}} \right)^{-1}$$

and for  $m_{\text{DM}} = 100 \text{ GeV}$  and weak  $g$ :


$$\sigma \sim g^4 / m_{\text{DM}}^2 \sim 3 \text{ pb} \cdot \text{cm/s}$$



\* Lee, Weinberg (1977)  
FERMILAB-PUB-77/41-THY

$n(n_0)$	DM number density (at equilibrium)
$\dot{R}/R$	expansion rate
$\langle\sigma v\rangle$	DM annihilation cross section x velocity
$\Omega_X h^2$	physical X density

# TRANSMISSION



•AMS



TDRS Satellites



Astronaut at ISS AMS Laptop

**Ku-Band**  
High Rate (down):  
Events <10Mbit/s

**S-Band**  
Low Rate (up & down):  
Commanding: 1 Kbit/s  
Monitoring: 30 Kbit/s



AMS Payload Operations Control and  
Science Operations Centers  
(POCC, SOC) at CERN since June 2011





AMS Computers  
at MSFC, AL






White Sands Ground  
Terminal, NM

# DÉTECTEUR

	$e^-$	P	He, Li, Be, ... Fe	$\gamma$	$e^+$	$\bar{P}, \bar{D}$	$\bar{He}, \bar{C}$
<b>TRD</b>							
<b>TOF</b>							
<b>Tracker</b>							
<b>RICH</b>							
<b>ECAL</b>							
<b>Physics example</b>	<b>Cosmic Ray Physics</b>				<b>Dark matter</b>		<b>Antimatter</b>


# POSITRON FRACTION

- **Key detectors** for this measurement
  - **TRD**
  - **Tracker**
    - $E/p$  close to 1 for electrons/positrons
  - **Calorimeter**
    - Based on 3D shower shape
- **Methodology**
  - **Selection** using the calorimeter variable
  - **Count** of  $e^+$  ( $Z > 0$ ) and  $e^-$  ( $Z < 0$ ) from a 2D fit on the TRD and tracker variables
  - Count for each **energy range**



# POSITRON FRACTION


- **Counts** of leptons after the selection
  - $Z > 0$  : count of **positrons**



- $Z < 0$  : count of **electrons**


# CHARGE CONFUSION

- For some energy range, difficulty to measure the **sign of the charge**  
⇒ confusion
- **Two sources**
  - Finite resolution of the tracker and multiple scattering
  - Production of secondary tracks along the path of the primary track



# COMBINED FLUX

- electron + positron flux measurement
  - Independent from **charge sign** measurement
  - **High selection efficiency** (70% at 1 TeV)



# MINIMAL MODEL

- Fit of the AMS data using a minimal model

- Positrons

- Secondary production
- + source

$$\Phi_{e^+} = C_{e^+} E^{-\gamma_{e^+}} + C_s E^{-\gamma_s} e^{-E/E_s}$$

- Electrons

- Primary and secondary production
- + same source

$$\Phi_{e^-} = C_{e^-} E^{-\gamma_{e^-}} + C_s E^{-\gamma_s} e^{-E/E_s}$$

- Simultaneous fit to

- Positron fraction from 2 GeV
- Combined flux from 2 GeV

# PROPAGATION

$$\frac{\partial \psi}{\partial t} - \nabla \cdot \{K(E) \nabla \psi\} - \frac{\partial}{\partial E} \{b(E) \psi\} = q(\mathbf{x}, t, E)$$

$$\psi = dn/dE$$

$$K(E) = K_0 \beta (\mathcal{R}/1 \text{ GV})^\delta$$

$$b(E) = \frac{E_0}{\tau_E} \epsilon^2 \quad \epsilon = E/E_0$$



$$q_{e^+}^{\text{DM}}(\mathbf{x}_S, E_S) = \frac{1}{2} \langle \sigma v \rangle \left\{ \frac{\rho_\chi(\mathbf{x}_S)}{m_\chi} \right\}^2 \left\{ g(E_S) \equiv \sum_i B_i \left. \frac{dN_{e^+}}{dE_S} \right|_i \right\}$$

$$g(E) = Q_0 \left( \frac{E_0}{E} \right)^\gamma \exp(-E/E_C)$$



$$\int_{E_{\min}}^{+\infty} E_S g(E_S) dE_S = fW_0.$$

# MINIMAL MODEL


Result from the fits



Prediction from the fits



Fits are satisfactory, which shows that the data can be described by a **common  $e^+/e^-$  source**



# The CR hadronic sector puzzle (*theoretical interpretation*)

Broken power law interpretations include:

- **diffusion effects** (source spectra assumed to be single power law):
  - non factorizable spatial and rigidity dependence of diffusion coefficient [N. Tomassetti, *Astrophys. J.* 752 , L13]
  - non linear diffusion on external turbulence (**self-generated waves**) above (**below**) the break [Blasi,Amato,Serpico, *PRL* 109]
- **acceleration effects** (observed features are imprinted on production spectra):
  - DSA acceleration non-linear effects (CR feed-back) [V. Ptuskin, V. Zirakashvili and E. S. Seo, *Astrophys. J.* 763]
  - Acceleration by different sources (e.g.: OB associations, SuperBubbles, W-R stars) [TStanev, Biermann & Gaisser, *Astron. Astrophys.* 274 , 902]
  - Weak re-acceleration [E. Seo and V. Ptuskin, *Astrophys. J.* 431]
- **local sources:**
  - Young nearby objects accounting for He harder spectrum are in tension with anisotropy measurements [Blasi, Amato, *JCAP* 1201 , 011]

Violation of universality of spectral indices interpretations include:

- e.g.: He accelerated “earlier” (with higher Mach number than proton) ?
  - He more efficient at injection than proton + slower decline with Mach number [Malkov, Diamond & Sagdeev, *Phys. Rev. Lett.* 108]
  - Variable He/p ion concentration in the medium swept by shocks [L. O. Drury, *Mon. Not. Roy. Astron. Soc.* 415 , 1807]

# ATLAS SUSY Searches\* - 95% CL Lower Limits

Status: EPS 2013

ATLAS Preliminary

$$\int \mathcal{L} dt = (4.4 - 22.9) \text{ fb}^{-1} \quad \sqrt{s} = 7, 8 \text{ TeV}$$

Model	$e, \mu, \tau, \gamma$	Jets	$E_T^{\text{miss}}$	$\int \mathcal{L} dt [\text{fb}^{-1}]$	Mass limit	Reference	
Inclusive Searches	MSUGRA/CMSSM	0	2-6 jets	Yes	20.3	$\tilde{q}, \tilde{g}$ 1.7 TeV	$m(\tilde{q})=m(\tilde{g})$
	MSUGRA/CMSSM	1 $e, \mu$	3-6 jets	Yes	20.3	$\tilde{g}$ 1.2 TeV	any $m(\tilde{q})$
	MSUGRA/CMSSM	0	7-10 jets	Yes	20.3	$\tilde{g}$ 1.1 TeV	any $m(\tilde{q})$
	$\tilde{q}\tilde{q}, \tilde{q} \rightarrow q\tilde{\chi}_1^0$	0	2-6 jets	Yes	20.3	$\tilde{q}$ 740 GeV	$m(\tilde{\chi}_1^0)=0 \text{ GeV}$
	$\tilde{g}\tilde{g}, \tilde{g} \rightarrow q\tilde{q}\tilde{\chi}_1^0$	0	2-6 jets	Yes	20.3	$\tilde{g}$ 1.3 TeV	$m(\tilde{\chi}_1^0)=0 \text{ GeV}$
	$\tilde{g}\tilde{g}, \tilde{g} \rightarrow q\tilde{q}\tilde{\chi}_1^0 \rightarrow qqW \pm \tilde{\chi}_1^0$	1 $e, \mu$	3-6 jets	Yes	20.3	$\tilde{g}$ 1.18 TeV	$m(\tilde{\chi}_1^0) < 200 \text{ GeV}, m(\tilde{\chi}^{\pm}) = 0.5(m(\tilde{\chi}_1^0) + m(\tilde{g}))$
	$\tilde{g}\tilde{g} \rightarrow qq\tilde{q}\tilde{\ell}(\tilde{\ell})\tilde{\chi}_1^0\tilde{\chi}_1^0$	2 $e, \mu$ (SS)	3 jets	Yes	20.7	$\tilde{g}$ 1.1 TeV	$m(\tilde{\chi}_1^0) < 650 \text{ GeV}$
	GMSB ( $\tilde{\ell}$ NLSP)	2 $e, \mu$	2-4 jets	Yes	4.7	$\tilde{g}$ 1.24 TeV	$\tan\beta < 15$
	GMSB ( $\tilde{\tau}$ NLSP)	1-2 $\tau$	0-2 jets	Yes	20.7	$\tilde{g}$ 1.4 TeV	$\tan\beta > 18$
	GGM (bino NLSP)	2 $\gamma$	0	Yes	4.8	$\tilde{g}$ 1.07 TeV	$m(\tilde{\chi}_1^0) > 50 \text{ GeV}$
	GGM (wino NLSP)	1 $e, \mu + \gamma$	0	Yes	4.8	$\tilde{g}$ 619 GeV	$m(\tilde{\chi}_1^0) > 50 \text{ GeV}$
	GGM (higgsino-bino NLSP)	$\gamma$	1 $b$	Yes	4.8	$\tilde{g}$ 900 GeV	$m(\tilde{\chi}_1^0) > 220 \text{ GeV}$
	GGM (higgsino NLSP)	2 $e, \mu$ (Z)	0-3 jets	Yes	5.8	$\tilde{g}$ 690 GeV	$m(H) > 200 \text{ GeV}$
Gravitino LSP	0	mono-jet	Yes	10.5	$F^{1/2}$ scale 645 GeV	$m(\tilde{g}) > 10^{-4} \text{ eV}$	
3 <sup>rd</sup> gen. $\tilde{g}$ med.	$\tilde{g} \rightarrow b\tilde{b}\tilde{\chi}_1^0$	0	3 $b$	Yes	20.1	$\tilde{g}$ 1.2 TeV	$m(\tilde{\chi}_1^0) < 600 \text{ GeV}$
	$\tilde{g} \rightarrow t\tilde{t}\tilde{\chi}_1^0$	0	7-10 jets	Yes	20.3	$\tilde{g}$ 1.14 TeV	$m(\tilde{\chi}_1^0) < 200 \text{ GeV}$
	$\tilde{g} \rightarrow t\tilde{t}\tilde{\chi}_1^0$	0-1 $e, \mu$	3 $b$	Yes	20.1	$\tilde{g}$ 1.34 TeV	$m(\tilde{\chi}_1^0) < 400 \text{ GeV}$
	$\tilde{g} \rightarrow b\tilde{t}\tilde{\chi}_1^0$	0-1 $e, \mu$	3 $b$	Yes	20.1	$\tilde{g}$ 1.3 TeV	$m(\tilde{\chi}_1^0) < 300 \text{ GeV}$
3 <sup>rd</sup> gen. squarks direct production	$\tilde{b}_1\tilde{b}_1, \tilde{b}_1 \rightarrow b\tilde{\chi}_1^0$	0	2 $b$	Yes	20.1	$\tilde{b}_1$ 100-630 GeV	$m(\tilde{\chi}_1^0) < 100 \text{ GeV}$
	$\tilde{b}_1\tilde{b}_1, \tilde{b}_1 \rightarrow t\tilde{\chi}_1^0$	2 $e, \mu$ (SS)	0-3 $b$	Yes	20.7	$\tilde{b}_1$ 430 GeV	$m(\tilde{\chi}_1^0) < 2 m(\tilde{\chi}_1^0)$
	$\tilde{t}_1\tilde{t}_1$ (light), $\tilde{t}_1 \rightarrow b\tilde{\chi}_1^0$	1-2 $e, \mu$	1-2 $b$	Yes	4.7	$\tilde{t}_1$ 167 GeV	$m(\tilde{\chi}_1^0) = 55 \text{ GeV}$
	$\tilde{t}_1\tilde{t}_1$ (light), $\tilde{t}_1 \rightarrow Wb\tilde{\chi}_1^0$	2 $e, \mu$	0-2 jets	Yes	20.3	$\tilde{t}_1$ 220 GeV	$m(\tilde{\chi}_1^0) = m(\tilde{t}_1) - m(W) - 50 \text{ GeV}, m(\tilde{t}_1) < m(\tilde{\chi}_1^0)$
	$\tilde{t}_1\tilde{t}_1$ (medium), $\tilde{t}_1 \rightarrow t\tilde{\chi}_1^0$	2 $e, \mu$	2 jets	Yes	20.3	$\tilde{t}_1$ 225-525 GeV	$m(\tilde{\chi}_1^0) < 0 \text{ GeV}$
	$\tilde{t}_1\tilde{t}_1$ (medium), $\tilde{t}_1 \rightarrow b\tilde{\chi}_1^0$	0	2 $b$	Yes	20.1	$\tilde{t}_1$ 150-580 GeV	$m(\tilde{\chi}_1^0) < 200 \text{ GeV}, m(\tilde{\chi}_1^{\pm}) - m(\tilde{\chi}_1^0) = 5 \text{ GeV}$
	$\tilde{t}_1\tilde{t}_1$ (heavy), $\tilde{t}_1 \rightarrow t\tilde{\chi}_1^0$	1 $e, \mu$	1 $b$	Yes	20.7	$\tilde{t}_1$ 200-610 GeV	$m(\tilde{\chi}_1^0) = 0 \text{ GeV}$
	$\tilde{t}_1\tilde{t}_1$ (heavy), $\tilde{t}_1 \rightarrow t\tilde{\chi}_1^0$	0	2 $b$	Yes	20.5	$\tilde{t}_1$ 320-660 GeV	$m(\tilde{\chi}_1^0) = 0 \text{ GeV}$
	$\tilde{t}_1\tilde{t}_1, \tilde{t}_1 \rightarrow c\tilde{\chi}_1^0$	0	mono-jet $c$ -tag	Yes	20.3	$\tilde{t}_1$ 200 GeV	$m(\tilde{t}_1) - m(\tilde{\chi}_1^0) < 85 \text{ GeV}$
	$\tilde{t}_1\tilde{t}_1$ (natural GMSB)	2 $e, \mu$ (Z)	1 $b$	Yes	20.7	$\tilde{t}_1$ 500 GeV	$m(\tilde{\chi}_1^0) > 150 \text{ GeV}$
$\tilde{t}_2\tilde{t}_2, \tilde{t}_2 \rightarrow \tilde{t}_1 + Z$	3 $e, \mu$ (Z)	1 $b$	Yes	20.7	$\tilde{t}_2$ 520 GeV	$m(\tilde{t}_1) = m(\tilde{\chi}_1^0) + 180 \text{ GeV}$	
EW direct	$\tilde{\ell}_L\tilde{\ell}_L, \tilde{\ell}_L \rightarrow \tilde{\ell}\tilde{\chi}_1^0$	2 $e, \mu$	0	Yes	20.3	$\tilde{\ell}$ 85-315 GeV	$m(\tilde{\chi}_1^0) = 0 \text{ GeV}$
	$\tilde{\chi}_1^{\pm}\tilde{\chi}_1^{\pm}, \tilde{\chi}_1^{\pm} \rightarrow \tilde{\ell}\nu(\tilde{\nu})$	2 $e, \mu$	0	Yes	20.3	$\tilde{\chi}_1^{\pm}$ 125-450 GeV	$m(\tilde{\chi}_1^0) = 0 \text{ GeV}, m(\tilde{\ell}, \tilde{\nu}) = 0.5(m(\tilde{\chi}_1^{\pm}) + m(\tilde{\chi}_1^0))$
	$\tilde{\chi}_1^{\pm}\tilde{\chi}_1^{\pm}, \tilde{\chi}_1^{\pm} \rightarrow \tilde{\tau}\nu(\tilde{\nu})$	2 $\tau$	0	Yes	20.7	$\tilde{\chi}_1^{\pm}$ 180-330 GeV	$m(\tilde{\chi}_1^0) = 0 \text{ GeV}, m(\tilde{\tau}, \tilde{\nu}) = 0.5(m(\tilde{\chi}_1^{\pm}) + m(\tilde{\chi}_1^0))$
	$\tilde{\chi}_1^{\pm}\tilde{\chi}_1^{\pm} \rightarrow \tilde{\ell}_L\nu_L\tilde{\ell}(\tilde{\nu}_L)\nu(\tilde{\nu}_L)$	3 $e, \mu$	0	Yes	20.7	$\tilde{\chi}_1^{\pm}, \tilde{\chi}_2^0$ 600 GeV	$m(\tilde{\chi}_1^0) = m(\tilde{\chi}_2^0), m(\tilde{\chi}_1^0) = 0, m(\tilde{\ell}, \tilde{\nu}) = 0.5(m(\tilde{\chi}_1^{\pm}) + m(\tilde{\chi}_1^0))$
	$\tilde{\chi}_1^{\pm}\tilde{\chi}_2^0 \rightarrow W\tilde{\chi}_1^0\tilde{\chi}_2^0$	3 $e, \mu$	0	Yes	20.7	$\tilde{\chi}_1^{\pm}, \tilde{\chi}_2^0$ 315 GeV	$m(\tilde{\chi}_1^0) = m(\tilde{\chi}_2^0), m(\tilde{\chi}_1^0) = 0$ , sleptons decoupled
Long-lived particles	Direct $\tilde{\chi}_1^{\pm}\tilde{\chi}_1^{\pm}$ prod., long-lived $\tilde{\chi}_1^{\pm}$	Disapp. trk	1 jet	Yes	20.3	$\tilde{\chi}_1^{\pm}$ 270 GeV	$m(\tilde{\chi}_1^0) - m(\tilde{\chi}_1^{\pm}) = 160 \text{ MeV}, \tau(\tilde{\chi}_1^{\pm}) = 0.2 \text{ ns}$
	Stable, stopped $\tilde{g}$ R-hadron	0	1-5 jets	Yes	22.9	$\tilde{g}$ 857 GeV	$m(\tilde{\chi}_1^0) = 100 \text{ GeV}, 10 \mu\text{s} < \tau(\tilde{g}) < 1000 \text{ s}$
	GMSB, stable $\tilde{\tau}, \tilde{\chi}_1^0 \rightarrow \tilde{\tau}(\tilde{e}, \tilde{\mu}) + \tau(e, \mu)$	1-2 $\mu$	0	-	15.9	$\tilde{\chi}_1^0$ 475 GeV	$10 < \tan\beta < 50$
	GMSB, $\tilde{\chi}_1^0 \rightarrow \gamma G$ , long-lived $\tilde{\chi}_1^0$	2 $\gamma$	0	Yes	4.7	$\tilde{\chi}_1^0$ 230 GeV	$0.4 < \tau(\tilde{\chi}_1^0) < 2 \text{ ns}$
	$\tilde{\chi}_1^0 \rightarrow qq\mu$ (RPV)	1 $\mu$	0	Yes	4.4	$\tilde{q}$ 700 GeV	$1 \text{ mm} < c\tau < 1 \text{ m}, \tilde{g}$ decoupled
RPV	LFV $pp \rightarrow \tilde{\nu}_\tau + X, \tilde{\nu}_\tau \rightarrow e + \mu$	2 $e, \mu$	0	-	4.6	$\tilde{\nu}_\tau$ 1.61 TeV	$\lambda_{311} = 0.10, \lambda_{132} = 0.05$
	LFV $pp \rightarrow \tilde{\nu}_\tau + X, \tilde{\nu}_\tau \rightarrow e(\mu) + \tau$	1 $e, \mu + \tau$	0	-	4.6	$\tilde{\nu}_\tau$ 1.1 TeV	$\lambda_{311} = 0.10, \lambda_{1(2)33} = 0.05$
	Bilinear RPV CMSSM	1 $e, \mu$	7 jets	Yes	4.7	$\tilde{q}, \tilde{g}$ 1.2 TeV	$m(\tilde{q}) = m(\tilde{g}), c\tau_{LSP} < 1 \text{ mm}$
	$\tilde{\chi}_1^{\pm}\tilde{\chi}_1^{\pm}, \tilde{\chi}_1^{\pm} \rightarrow W\tilde{\chi}_1^0\tilde{\chi}_1^{\pm} \rightarrow ee\tilde{\nu}_\mu, e\mu\tilde{\nu}_e$	4 $e, \mu$	0	Yes	20.7	$\tilde{\chi}_1^{\pm}$ 760 GeV	$m(\tilde{\chi}_1^0) > 300 \text{ GeV}, \lambda_{121} > 0$
	$\tilde{\chi}_1^{\pm}\tilde{\chi}_1^{\pm}, \tilde{\chi}_1^{\pm} \rightarrow W\tilde{\chi}_1^0\tilde{\chi}_1^{\pm} \rightarrow \tau\tau\tilde{\nu}_e, e\tau\tilde{\nu}_\tau$	3 $e, \mu + \tau$	0	Yes	20.7	$\tilde{\chi}_1^{\pm}$ 350 GeV	$m(\tilde{\chi}_1^0) > 80 \text{ GeV}, \lambda_{133} > 0$
	$\tilde{g} \rightarrow qq\tilde{q}$	0	6 jets	-	4.6	$\tilde{g}$ 666 GeV	
$\tilde{g} \rightarrow \tilde{t}_1 t, \tilde{t}_1 \rightarrow bs$	2 $e, \mu$ (SS)	0-3 $b$	Yes	20.7	$\tilde{g}$ 880 GeV		
Other	Scalar gluon	0	4 jets	-	4.6	sgluon 100-287 GeV	incl. limit from 1110.2693
	WIMP interaction (D5, Dirac $\chi$ )	0	mono-jet	Yes	10.5	$M^*$ scale 704 GeV	$m(\chi) < 80 \text{ GeV}$ , limit of $c$ for D8

$\sqrt{s} = 7 \text{ TeV}$   
full data

$\sqrt{s} = 8 \text{ TeV}$   
partial data

$\sqrt{s} = 8 \text{ TeV}$   
full data

$10^{-1}$

1


Mass scale [TeV]

\*Only a selection of the available mass limits on new states or phenomena is shown. All limits quoted are observed minus  $1\sigma$  theoretical signal cross section uncertainty.

# Current limits: neutralino/chargino

7


canonical case



$m_{\tilde{\chi}_1^\pm} > 103.5$  GeV  
for  $m_{\text{snue}} > 300$  GeV

LEPSUSYWG/01-03.1


degenerate case



$m_{\tilde{\chi}_1^\pm} > 91.9 / 92.4$  GeV

LEPSUSYWG/02-04.1

$m_{\tilde{\chi}_1^0} > 47/50$  GeV  
(CMSSM, mSUGRA)  
No mass limit in general



7  
LEPSUSYWG/04-01.1

S. Su