



Summer School in **Particle and Astroparticle physics**  
of Annecy-le-Vieux

**20-26 July 2017**



## A computing exercise using ROOT

**Aim:** give a taste of data analysis @ LHC

- What is ROOT ?
  - ROOT is an object-oriented C++ analysis package
  - User-compiled code can be called to produce 1-d, 2-d, and 3-d graphics and histograms...



<https://root.cern.ch>

# Outline

---

- Kinematic variables used in the analysis of  $p - p$  collisions
- Useful relations
- Concept of invariant mass: example: ‘inclusive’ Z boson production
- Kinematics of  $p - p$  collisions
  
- Analysis in  $p - p$  collisions :
  - \* Signal: Production of a W and a Z  $p - p \rightarrow WZ X$
  - \* Background: Production of a pair of top-antitop
- Example: Macro.C

In all the following slides we assume the speed of the light

$$c=1$$

# Variables used in the analysis of $p$ - $p$ collisions

A particle ( Z, W, e+, e-, etc ...) is described by its **four-momentum**:

$$\tilde{p} = (E, p_x, p_y, p_z)$$

The particle mass is  $m = \sqrt{E^2 - p_x^2 - p_y^2 - p_z^2}$

When dealing with  $p$ - $p$  collision the following variables are used:

For each particle ( Z, W, e+, e-, etc ...):

- 1. Transverse momentum/energy :  $p_T = p \sin \theta$        $E_T = E \sin \theta$

- 2. Rapidity

or Pseudorapidity


$$Y = \frac{1}{2} \ln \frac{E + p_z}{E - p_z}$$

$$\eta = -\ln \left( \tan \frac{\theta}{2} \right)$$

- 3. Azimuthal angle


$$\Phi$$

Why?



# Variables used in the analysis of $p - p$ collisions

Why  $p_T, Y$ ? Many reasons.



1. The longitudinal momentum of initial partons is unknown while we know that  $\vec{p}_T^{\text{initial partons}} \sim 0$

→ To exploit momentum conservation

use transverse quantities (in the plane  $\perp$  to the beam) →  $p_T$


2.  $p_T$  and  $\Delta Y$  are invariants for Lorentz transformations along the z axis
3.  $\sum_{\text{initial partons}} \vec{p}_T = \sum_{\text{vis}} \vec{p}_T + \sum_{\text{invis}} \vec{p}_T \approx 0$  → Allows to evaluate the  $p_T$  of particles not detected (v)

$$\sum_{\text{invis}} \vec{p}_T = - \sum_{\text{vis}} \vec{p}_T \quad |\sum_{\text{invis}} \vec{p}_T| \text{ is the "missing } E_T \text{"}$$

4. The “interesting” physics is due to hard scattering processes → high  $p_T$  particles (selection of high  $p_T$  particles assures “interesting” physics)

## Useful relations

---



$$p_x = p_T * \cos(\Phi);$$

$$p_y = p_T * \sin(\Phi);$$

$$p_z = E * \tanh(\eta);$$

$$\begin{aligned} p_T &= p \sin \theta \\ \eta &= -\ln (\tan \frac{\theta}{2}) \end{aligned}$$

- $m \ll E \rightarrow Y \approx \eta$  ( $\eta$  doesn't require particle identification)
- $m \ll E \rightarrow p_T \approx E_T \quad E_T = E \sin \theta$

## Concept of invariant mass: inclusive Z boson production

$p - p \rightarrow Z X$

With  $Z \rightarrow e+e-$   
 $X = p_1, p_2, p_3, \dots$

Very 'clean' processes (low bkg)!!

Invariant mass  $M_{ee}$  of  $ee$  system from the  
 4-momentum conservation  
 ( it allows to measure the Z mass,  $M_Z$  ):


$$\tilde{p}_Z^2 = (\tilde{p}_{e1} + \tilde{p}_{e2})^2$$

$$M_{ee}^2 = (\tilde{p}_{e1} + \tilde{p}_{e2})^2 \approx 2(E_{e1}E_{e2} - |\vec{p}_{e1}| |\vec{p}_{e2}| \cos \vartheta_{e1 e2})$$

$$M_{ee} \approx \sqrt{2} E_{e1} E_{e2} (1 - \cos \vartheta_{e1 e2})$$

(the electron mass is neglected)

Why  $M_{ee}$  gives a distribution  
 and not a single value?



1.  $\Delta E * \Delta t > \hbar/2$     $\Delta m * \tau > \hbar/2$

$$\Gamma * \tau > \hbar/2$$


width

lifetime

2. Experimental resolution

# Our signal : Production of a W and a Z

$p-p$  'hard' collisions in the  $q_1 \bar{q}_2$  center of mass:




## Kinematics of $p-p$ collisions

\* 4-mom of the initial partons :  $[ (x_1+x_2)E_{beam}, 0, 0, (x_1-x_2) p_{beam}]$

$$0 < x_{1,2} < 1$$

# Our background: Production of a pair of top-antitop

$p-p$  'hard' collisions. In the  $q_1 \bar{q}_2$  center of mass:



The **b quarks** flies few cm before desintegrating  $\rightarrow e^+(\mu^+) \text{ from } \mathbf{b \ quark}$  do not emerge from the primary vertex  $\rightarrow$  Higher distance from the primary vertex (higher "impact parameter")

## Aim of the exercise:

- 1) look at some important variables,
- 2) build the Z invariant mass,
- 3) how one can discriminate between the 'signal' and the 'background'

**GRASPA2017explanation.pptx.pdf (this slides)**

**Exercise2017.pdf** (what we ask to do)  
**macro.C** (draft of analysis program)  
**Selected\_All\_EEM.root** (« data » (simulated data))  
**macro\_final.C** (solution: final analysis program)  
**Tutorial\_ROOT\_Bose.pdf** (an old by simple Root manual)

# 1) an **input file** containing the physics: **Selected\_All\_EEM.root**

```
==== MOST ENERGETIC LEPTON FROM THE Z  
Br 4 :pt1 : pt1  
Br 5 :etal1 : eta1  
Br 6 :phi1 : phi1  
Br 7 :E1 : E1  
  
==== SECOND ENERGETIC LEPTON FROM THE Z  
Br 8 :pt2 : pt2  
Br 9 :eta2 : eta2  
Br 10 :phi2 : phi2  
Br 11 :E2 : E2  
  
==== LEPTON FROM W  
Br 12 :pt3 : pt3  
Br 13 :eta3 : eta3  
Br 14 :phi3 : phi3  
Br 15 :E3 : E3
```

**List of variables given per each collision event (kinematics of the final state leptons)**

## 2) Instructions to make the computing exercise : **Exercise2017.pdf**



### COMPUTING EXERCISE Study of the production of a pair of gauge bosons ( $W$ and $Z$ ) at the LHC

The data to analyse are organised into a 'Root n-tuple' which we will provide to you. The Root n-tuple is a file containing information about the kinematics of "events", each resulting from a proton-proton interaction.

These events have three leptons (electrons or muons) and are of two kinds:

### 3) A skeleton of an analysis program using ROOT: macro.C

```
#include "TCanvas.h"
#include "TROOT.h"
#include "TFile.h"
#include "TTree.h"
#include "TBrowser.h"
#include "TH2.h"
#include "TRandom.h"

void tree1r()
{
    // Read Selected_All_EEM.root file
    //Root file
    TFile *f = new TFile("Selected_All_EEM.root");

    // Signal events
    TTree *sig = (TTree*)f->Get("WZSignal");
    Double_t pt1, eta1, phi1, E1;
    Double_t pt2, eta2, phi2, E2;
    Double_t pt3, eta3, phi3, E3;
    Double_t MZ, MET, trackd0cutWMu, TrackIsoWmu;
    Double_t Weight;

    //get some variables for SIGNAL EVENTS
    sig->SetBranchAddress("pt1",&pt1);
```

# Example of analysis program

macro.C

23/07/2013 00:21

```
#include "TCanvas.h"
#include "TROOT.h"
#include "TFile.h"
#include "TTree.h"
#include "TBrowser.h"
#include "TH2.h"
#include "TRandom.h"

void tree1r()
{
    // Read Selected_All_EEM.root file
    //Root file
    TFile *f = new TFile("Selected_All_EEM.root");

    // Signal events
    TTree *sig = (TTree*)f->Get("WZSignal");
    Double_t pt1, eta1, phi1, E1;
    Double_t pt2, eta2, phi2, E2;
    Double_t pt3, eta3, phi3, E3;
    Double_t MZ, MET, trackd0cutWMu, TrackIsoWMu;
    Double_t Weight;

    //get some variables for SIGNAL EVENTS
    sig->SetBranchAddress("pt1",&pt1);
    sig->SetBranchAddress("eta1",&eta1);
    sig->SetBranchAddress("phi1",&phi1);
    sig->SetBranchAddress("E1",&E1);
    sig->SetBranchAddress("MZ",&MZ);
    sig->SetBranchAddress("Weight",&Weight);
    // add other variables ...
```

**Header files**

**Open the input file**

**Access the Signal info**

**Define the name  
variables per each SIGNAL lepton**

```

////get some variables for BACKGROUND EVENTS
TTree *ttbar = (TTree*)f->Get("ttbar");
Double_t pt1_bkg, eta1_bkg, phi1_bkg, E1_bkg;
Double_t MZ_bkg;
Double_t Weight_bkg;

//get some variables for ttbar
ttbar->SetBranchAddress("pt1",&pt1_bkg);
ttbar->SetBranchAddress("eta1",&eta1_bkg);
ttbar->SetBranchAddress("phi1",&phi1_bkg);
ttbar->SetBranchAddress("E1",&E1_bkg);
ttbar->SetBranchAddress("MZ",&MZ_bkg);
ttbar->SetBranchAddress("Weight",&Weight_bkg);
// add other variables ...

//create two histograms (for sig and ttbar)
TH1F *h_MZ    = new TH1F("h_MZ","MZ distribution All events",40,65,115);
TH1F *h_MZ_bkg = new TH1F("h_MZ_bkg","MZ distribution BKG",40,65,115);
TH1F *h_MZ_sig = new TH1F("h_MZ_sig","MZ distribution SIG",40,65,115);

//read all SIGNAL entries and fill the histograms
Int_t nentries = (Int_t)sig->GetEntries();

for (Int_t i=0;i<nentries_bkg;i++) {
    ttbar->GetEntry(i);
    h_MZ_bkg->Fill(MZ_bkg,Weight_bkg);
    h_MZ->Fill(MZ_bkg,Weight);
}

```

**Access the background info**

**Define the name  
variables per each bkg lepton**

**Loop on events**

Page 1 of 2

```
// example how Draw and save histograms
TCanvas *c = new TCanvas();
c->cd();
h_MZ_sig->Draw();
h_MZ_bkg->SetLineColor(kRed);
h_MZ_bkg->Draw("same");

c->Print("test_MZ.eps");
}
```

```
void macro()
{
    tree1r();
}
```

**Draw and save histograms**

**Main program**

---

**To start root you may type:**

**root**

**root [1] .x macro.C**

**and look at what you get ....**

Useful links about root:

[https://root.cern.ch/root/html/doc/guides/primer\(ROOTPrimer.pdf](https://root.cern.ch/root/html/doc/guides/primer(ROOTPrimer.pdf)

## ROOT Tutorial

4) A 'manual' with ROOT instructions: [Tutorial\\_ROOT\\_Bose.pdf](#)

Tulika Bose  
Brown University  
NEPPSR 2007

<http://www.phys.vt.edu/~dayabay/Presentations/090916.dm.Root1.pdf>

**Have fun !!**



**W(jj) Z (jj)**

**jj = J (1 fat jet)**

## Another example: search for di-boson resonances

- Is there something hiding in the data, waiting to be discovered?

