



Course on Physics at the LHC

Lecture 1

Joao Varela



LABORATÓRIO DE INSTRUMENTAÇÃO
E FÍSICA EXPERIMENTAL DE PARTÍCULAS
partículas e tecnologia

Lisbon, PORTUGAL
06 MARCH - 09 MAY

A specialized course on the Physics at the Large Hadron Collider is organized by LIP in the framework of the International Doctorate Network in Particle Physics, Astrophysics and Cosmology (IDPASC).

The course is intended for under-graduate or graduate students with basic training in Particle Physics.

The objective of the Course is to introduce the physics, analysis methods and results on the physics areas covered by the LHC experiments.

- Experimental program at the LHC
- Standard Model at the LHC
- Detectors
- Statistics
- Top quark physics
- Higgs Physics
- Supersymmetry
- Exotic processes and Dark Matter
- Heavy flavor physics and rare decays searches
- Matter at high density and temperature



Prof. João Varela
Coordinator/lecturer



Dr. Michele Gallinaro
Coordinator/lecturer



Dr. Pedro Silva
Invited lecturer (CERN)



Dr. Pietro Vischia
Invited lecturer (Univ. Oviedo)



Prof. António Onofre
Resident lecturer



Dr. Ricardo Gonçalves
Resident lecturer



Dr. Patricia Muñio
Resident lecturer



Dr. Pedrame Bargassa
Resident lecturer



Dr. Nuno Leonardo
Resident lecturer




Prof. João Seixas
Resident lecturer

Background bibliography

Students are recommended to refresh their knowledge in particle physics prior to the Course. Suggested bibliography:

- F. Halzen and A.D.Martin, ' Quarks and Leptons ', John Wiley and Sons (1984)
- D. Griffiths, ' Introduction to Elementary Particles ', John Wiley and Sons (1987)
- B.R.Martin, G. Shaw, ' Particle Physics ', John Wiley and Sons (1999)
- D. Perkins, ' Introduction to High Energy Physics', Cambridge University Press (2000)

The LHC physics case



Particle physics is a modern name for the centuries old effort to understand the basic laws of physics.

Edward Witten

Aims to answer the two following questions:


What are the elementary constituents of matter ?

What are the forces that determine their behavior?

Experimentally

Get particles to interact and study what happens


Constituents of matter along History



The Standard Model

Over the last ~100 years: The combination of Quantum Field Theory and discovery of many particles has led to

- **The Standard Model of Particle Physics**
 - With a new “Periodic Table” of fundamental elements




One of the greatest achievements of 20th Century Science

$$L_H = \frac{1}{2}(\partial_\mu H)^2 - m_H^2 H^2 - h\lambda H^3 - \frac{h}{4}H^4 + \frac{g^2}{4}(W_\mu^+ W^\mu + \frac{1}{2\cos^2\theta_W} Z_\mu Z^\mu)(\lambda^2 + 2\lambda H + H^2) + \sum_{l,q,q'} (\frac{m_l}{\lambda} \bar{l}l + \frac{m_q}{\lambda} \bar{q}q + \frac{m_{q'}}{\lambda} \bar{q}'q')H$$

SM confirmed by data

	I	II	III	
mass →	2.4 MeV/c ²	1.27 GeV/c ²	171.2 GeV/c ²	0
charge →	$\frac{2}{3}$	$\frac{2}{3}$	$\frac{2}{3}$	0
spin →	$\frac{1}{2}$	$\frac{1}{2}$	$\frac{1}{2}$	1
name →	u up	c charm	t top	γ photon
Quarks	4.8 MeV/c ²	104 MeV/c ²	4.2 GeV/c ²	0
	$-\frac{1}{3}$	$-\frac{1}{3}$	$-\frac{1}{3}$	0
	$\frac{1}{2}$	$\frac{1}{2}$	$\frac{1}{2}$	1
	d down	s strange	b bottom	g gluon
Leptons	<2.2 eV/c ²	<0.17 MeV/c ²	<15.5 MeV/c ²	91.2 GeV/c ²
	0	0	0	0
	$\frac{1}{2}$	$\frac{1}{2}$	$\frac{1}{2}$	1
	ν_e electron neutrino	ν_μ muon neutrino	ν_τ tau neutrino	Z⁰ Z boson
Gauge bosons	0.511 MeV/c ²	105.7 MeV/c ²	1.777 GeV/c ²	80.4 GeV/c ²
	-1	-1	-1	±1
	$\frac{1}{2}$	$\frac{1}{2}$	$\frac{1}{2}$	1
	e electron	μ muon	τ tau	W[±] W boson

STANDARD MODEL OF ELEMENTARY PARTICLES



Confirmed at sub 1% level!

$$\mathcal{L} = -\frac{1}{4} F_{\mu\nu} F^{\mu\nu} + i\bar{\psi}\not{D}\psi + \text{h.c.}$$

In the simplest model the interactions are symmetrical and particles do not have mass

The symmetry between the electromagnetic and the weak interactions is broken:

- Photon do not have mass
- W, Z do have a mass $\sim 80\text{-}90\text{ GeV}$

Higgs mechanism:

mass of W and Z results from the interactions with the Higgs field

The Standard Model would fail at high energy without the Higgs particle or other ‘new physics’

Based on the available data and on quite general theoretical insights it was expected that the ‘**new physics**’ would manifest at an energy around

1 Tera-electronVolt = 10^{12} electronVolt

accessible at the LHC for the first time



Beyond the standard model

The Standard Model answers many of the questions about the structure of matter. But the Standard Model is not complete; there are still many unanswered questions.

Why do we observe matter and almost no antimatter if we believe there is a symmetry between the two in the universe?


What is this "dark matter" that we can't see that has visible gravitational effects in the cosmos?

Are quarks and leptons actually fundamental, or made up of even more fundamental particles?

Why are there three generations of quarks and leptons? What is the explanation for the observed pattern for particle masses?

How does gravity fit into all of this?

Forces and expansion of the Universe




The dark side of the Universe

Long standing problem:

We know that ordinary matter is only $\sim 4\%$ of the matter-energy in the Universe.

What is the remaining 96%?



The LHC may help to solve this problem, discovering **dark matter**


The Universe expansion is accelerating

In 1998, two groups used distant **Supernovae** to measure the expansion rate of the universe: Perlmutter et al. (Supernova Cosmology Project), and Schmidt et al. (High-z Supernova Team)

They got the same result:

The Universe expansion is accelerating


Some form of energy (dark energy) fills space



Cosmological inflation

In the very early universe space undergoes a dramatic exponential expansion.

Explains why the Universe has a uniform Temperature (3 K) and why space-time has a flat geometry




The inflation theory was developed independently in the late 1970's by Alan Guth, Alexey Starobinsky, and others

Higgs and hierarchy problem

In the SM the Higgs mass is a huge problem:

- Virtual particles in quantum loops contribute to the Higgs mass
- Contributions grow with Λ (upper scale of validity of the SM)
- Λ could be huge – e.g. the Plank scale (10^{19} GeV)
- Miraculous cancelations are needed to keep the Higgs mass < 1 TeV



$$m_h^2 = (m_h^2)_0 - \frac{1}{16\pi^2} \lambda^2 \Lambda^2 + \dots,$$

This is known as the hierarchy problem

Many possible theories

There are a large number of models which predict new physics at the TeV scale accessible at the LHC:

- Supersymmetry (SUSY)
- Extra dimensions
- Extended Higgs Sector e.g. in SUSY Models
- Grand Unified Theories (SU(5), O(10), E6, ...)
- Leptoquarks
- New Heavy Gauge Bosons
- Compositeness


Any of this could still be found at the LHC

Supersymmetry

New fundamental symmetry:

- Every fermion should have a massive "shadow" boson
- Every boson should have a massive "shadow" fermion.
-

This relationship between fermions and bosons is called supersymmetry (SUSY)



Heavy versions of every quark and lepton

Supersymmetry is broken

Could DM be SUSY particles?

For every “normal” force quanta (boson), there are supersymmetric partners:

photon

W, Z bosons

gluon

Higgs boson

photino

Wino, Zino


gluino

higgsino


These “...inos” are prime suspects to be the galactic dark matter!

Relics from the Big Bang!

The temptation unification



SUSY and the Higgs mass



$$m_h^2 = (m_h^2)_0 - \frac{1}{16\pi^2} \lambda^2 \Lambda^2 + \dots,$$

Higgs mass:

- correction has quadratic divergence!
 - Λ a cut-off scale – e.g. Planck scale

Superpartners fix this:

- Need superpartners at mass $\sim 1\text{-}2$ TeV
 - Otherwise the logarithmic term becomes too large, which would require more fine-tuning.

Cancellation

$$m_h^2 = (m_h^2)_0 - \frac{1}{16\pi^2} \lambda^2 \Lambda^2 + \frac{1}{16\pi^2} \lambda^2 \Lambda^2 + \dots$$

$$\approx (m_h^2)_0 + \frac{1}{16\pi^2} (m_{\tilde{f}}^2 - m_f^2) \ln(\Lambda / m_{\tilde{f}}),$$

Extra dimensions


Space-time could have more than three space dimensions. The extra dimensions could be very small and undetected until now.

How can there be extra, smaller dimensions?

The acrobat can move forward and backward along the rope: **one dimension**

The flea can move forward and backward as well as side to side: **two dimensions**

But one of these dimensions is a small closed loop.




Understanding the Universe

History of the Universe

Accelerators

Telescopes

LHC – Energy Frontier




Particle Data Group, LBNL, © 2008. Supported by DOE and NSF


The LHC proton collider



Accelerator and Experiments




Accelerator and experiments layout




**Tiny bunches of counter-circulating protons.
Colliding head-on 40 million times each second.**


Collisions at LHC



Superconducting magnetic dipole





In the tunnel



It's empty!

Air pressure inside the two 27Km-long vacuum pipes (10^{-13} atm) is lower than on the moon.




It's cold!


27 Km of magnets are kept at 1.9 °K, colder than outer space, using over 100 tons of liquid helium.



In a *tiny* volume, temperatures one billion times hotter than the center of the sun.



The Experiments




General purpose LHC experiments

Advanced detectors comprising many layers, each designed to perform a specific task.

Together these layers allow to identify and precisely measure the energies of all stable particles produced in collisions.

Photons,
Electrons,
Muons,
Quarks
(as jets of particles)
Neutrinos
(as missing energy)



CMS DETECTOR

Total weight : 14,000 tonnes
Overall diameter : 15.0 m
Overall length : 28.7 m
Magnetic field : 3.8 T

STEEL RETURN YOKE
12,500 tonnes

SILICON TRACKERS

Pixel ($100 \times 150 \mu\text{m}$) ~16m²~66M channels
Microstrips ($80 \times 180 \mu\text{m}$) ~200m²~9.6M channels

CMS Detector

SUPERCONDUCTING SOLENOID

Niobium titanium coil carrying ~18,000A

MUON CHAMBERS

Barrel: 250 Drift Tube, 480 Resistive Plate Chambers
Endcaps: 468 Cathode Strip, 432 Resistive Plate Chambers

PRESHOWER


Silicon strips ~16m²~137,000 channels

FORWARD CALORIMETER


Steel + Quartz fibres ~2,000 Channels

CRYSTAL
ELECTROMAGNETIC
CALORIMETER (ECAL)
~76,000 scintillating PbWO₄ crystals


HADRON CALORIMETER (HCAL)
Brass + Plastic scintillator ~7,000 channels



ATLAS detectors



Detection of hadrons, e^\pm , γ and μ^\pm





1993-2008: detector R&D and construction



15 years !



Superconductor solenoid at 3.8 Tesla




ATLAS Toroidal System




Silicon Tracker

Diameter 2.4m
Length 5.4m
Volume 24.4m³
Running temperature -10⁰C
Dry atmosphere for 10 years






Used to reconstruct the trajectories of thousands of charge particles produced in the collisions

214m² silicon sensors
11.4 million silicon strips
65.9 million silicon pixels



ECAL Electromagnetic Calorimeter



Design Goal: Measure the energies of photons from a decay of the Higgs boson to precision of $\leq 0.5\%$

Parameter	Barrel	Endcaps
# of crystals	61200	14648
Volume	8.14m ³	2.7m ³
Xtal mass (t)	67.4	22.0

HCAL Hadronic Calorimeter

Detection of hadrons:


- protons, neutrons, pions, etc.

- CMS HCAL has three components:



- Barrel HCAL (HB)
- Endcap HCAL (HE)
- Forward HCAL (HF)

- Plastic scintillator and brass

- Quartz fibers and steel



Muon detectors






Drift Tubes (DT)
Cathode Strip Chambers (CSC)
Resistive Plate Chambers (RPC)




Trigger and readout electronics

Underground caverns




Electronics systems

Electronics systems in the Service Cavern.
About 150 racks occupy two floors.
Most electronics was designed and built
specifically for the experiment





Sep 2008: CMS detector ready for beams



Detector simulation

Simulation of proton-proton collision
making two dark matter particles



The LHC Computing Grid


The Grid unites computing resources of particle physics institutions around the world

The **World Wide Web** (invented at CERN) provides seamless access to information that is stored in many millions of different geographical locations

The **Grid** is an infrastructure that provides seamless access to computing power and data storage capacity distributed over the globe



Experimental challenges



High collision rate

Luminosity:

$$L = 10^{34} \text{ cm}^{-2}\text{s}^{-1}$$

$$= 10^7 \text{ Hz/mb}$$

Cross section:

$$\sigma \approx 100 \text{ mb}$$


$$\rightarrow N = L\sigma \approx 1 \text{ GHz}$$

However:

Bunch crossing rate: 40 MHz


\therefore Interactions/crossing ~ 25

This is a
real challenge !




Bunch crossing frequency


- LHC has 3564 bunches (2835 filled with protons)
- Crossing rate is 40 MHz
- Distance between bunches: $27\text{km} / 3600 = 7.5\text{m}$
- Distance between bunches in time: $7.5\text{m} / c = 25\text{ns}$
- Proton-proton collision per bunch crossing: ~ 25





- Proton bunches have a cigar shape, about 5 cm long and 20 microns diameter
- Each bunch has $1.5 \cdot 10^{11}$ protons
- At each crossing of bunches, about 25 collision occur
- The particles produced ($30 \times 25 = 750$ charged particles) are “seen” by the detector as a single image (event)



High radiation levels



Two-level trigger



Trigger system decide if the event is interesting to be recorded

Two-step process:

- **Level 1**: dedicated hardware processors
- **High level**: computer farm

Triggers and event selection

- Select processes that produce particles with high transverse energy
- Examples at $5 \times 10^{33} \text{ cm}^{-2}\text{s}^{-1}$
 - Single lepton and photon triggers ($P_T \sim 30 \text{ GeV}$)
 - Multiple lepton and photon triggers ($P_T \sim 15 \text{ GeV}$)
 - Missing transverse energy ($P_T \sim 50\text{-}100 \text{ GeV}$)
 - Multiple jet triggers ($P_T \sim 50\text{-}100 \text{ GeV}$)
- About 100 trigger conditions in L1 trigger table
- About 400 trigger conditions in HLT trigger table

LHC Page 1: stable beams

LHC Page1

Fill: 1729

E: 3500 GeV

22-04-2011 00:02:58

PROTON PHYSICS: STABLE BEAMS

Energy:

3500 GeV

I(B1):


5.50e+13

I(B2):

5.54e+13


FBCT Intensity and Beam Energy

Updated: 00:02:58



Instantaneous Luminosity

Updated: 00:02:54



Comments 22-04-2011 00:02:52 :

**** Stable Beams ****

World record for luminosity for
hadron machine
Automatic LUMI LEVELING in IP8

BIS status and SMP flags

B1 B2

Link Status of Beam Permits

true true

Global Beam Permit

true true

Setup Beam

false false

Beam Presence

true true

Moveable Devices Allowed In

true true

Stable Beams

true true

AFS: 50ns_480b+1small_424_12_468_36bpl15nj

PM Status B1

ENABLED

PM Status B2

ENABLED

March 30, 2010: CMS Page 1




2009: First p-p collisions at LHC

November 23, 2009
First collisions at 900 GeV




December 14, 2009
First collisions at 2.36 TeV

March 30, 2010
First collisions at 7 TeV



Tracking: secondary vertices

Basic variables relevant for B-tagging are well described by the simulation







Secondary vertices compatible with heavy flavor production




Photons and electrons

EM cluster energy




Jets and missing energy




ATLAS


Missing Transverse Energy








Di-jet mass




CMS



Rediscovery of resonances



Rediscovery of the Standard Model at LHC



End of Lecture 1